

COMPARATIVE STUDY REGARDING THE THERAPY OF TRICHINOSIS WITH THIABENDAZOLE AND MEBENDAZOLE, IN 250 CASES OF TRICHINELLOSIS CONFIRMED BY MEANS OF IMMUNOASSAY AND MUSCULAR BIOPSY

*L.Negruțiu, I.Iacobiciu, Rodica Potcoavă-Vasiliu, Narcisa Nicolescu,
L.Sălăgean, I.Marincu*

Clinic of Infectious Diseases; Department of Parasitology,
University of Medicine and Pharmacy, Timișoara;
"I.C.Cantacuzino" Institute, Bucharest

Our study compares the therapeutic efficiency of antiparasitic therapy with Thiabendazole and Mebendazole, respectively, in two groups of 125 patients with trichinellosis hospitalized in an interval of 4 years in the Clinic of Infectious Diseases Timișoara (1989-1994). The etiological diagnosis was established by direct visualisation of the parasite in muscular biopsy specimens and by means of ELISA and RPL (90 cases investigated at the Parasitology Laboratory of the I.C.Cantacuzino Institute - Bucharest). The clinical and Biological monitoring of some evolutive parameters permits a comparative study of the therapeutical results obtained with the two drugs in our patients.